

---

**CHEMISTRY**

**9701/51**

Paper 5 Planning, Analysis and Evaluation

**May/June 2018**

MARK SCHEME

Maximum Mark: 30

---

**Published**

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the May/June 2018 series for most Cambridge IGCSE™, Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

---

IGCSE™ is a registered trademark.

This document consists of **7** printed pages.

**PUBLISHED****Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

**GENERIC MARKING PRINCIPLE 1:**

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

**GENERIC MARKING PRINCIPLE 2:**

Marks awarded are always **whole marks** (not half marks, or other fractions).

**GENERIC MARKING PRINCIPLE 3:**

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

**GENERIC MARKING PRINCIPLE 4:**

Rules must be applied consistently e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

**GENERIC MARKING PRINCIPLE 5:**


Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

**GENERIC MARKING PRINCIPLE 6:**

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

**PUBLISHED**

<b>Question</b>	<b>Answer</b>	<b>Marks</b>
1(a)	Complete circuit with ammeter in series and DC power supply	<b>1</b>
	Anode, cathode and solution labelled	<b>1</b>
1(b)	wear gloves	<b>1</b>
	do not dispose into the water waste / sink <b>OR</b> do not put down drain / sewage <b>OR</b> put in waste bottles	<b>1</b>
1(c)	Mass (of electrode) before and after experiment <b>AND</b> mass unit	<b>1</b>
1(d)	charge = $0.5 \times 30 \times 60 = 900$ C	<b>1</b>
1(e)	$0.282 / 63.5 = 4.44 \times 10^{-3}$ (mol) <b>OR</b> 0.00444	<b>1</b>
1(f)	$(900 / 4.44 \times 10^{-3}) = 202702.7027$ C	<b>1</b>
1(g)	2 moles of electrons are <b>produced</b> / removed / released (so 2 Faradays <b>OR</b> $2 \times 96\,500$ )	<b>1</b>
1(h)	(Faraday) value is smaller <b>AND</b> (apparent) mass / moles / amount is more (for same charge passed)	<b>1</b>
1(i)	CuO is formed / oxidation of copper / carbon / soot is formed	<b>1</b>
1(j)	Some copper falls off the electrode during electrolysis / falls to the bottom of the beaker <b>OR</b> Some copper is lost during washing	<b>1</b>

Question	Answer	Marks
2(a)	Water bath/beaker of water containing thermometer around flask	1
	Controlled heat source or heater/temperature regulator	1
2(b)(i)	Moles $N_2 = 72 / 24\,000 = 0.003$ moles (1 mol $C_6H_5N_2^+Cl^- \rightarrow 1$ mol $N_2$ )	1
	Moles $C_6H_5N_2^+$ in $1000\text{ cm}^3$ solution = $0.003 \times (1000 / 200) = 1.50 \times 10^{-2}$ (mol)	1
2(b)(ii)	 <p style="text-align: center;">OR</p>	
	Axes (label with quantity or correct unit) and values correct	1
	Straight line from axis marks <b>OR</b> from 0,0 over most of the axes	1

**PUBLISHED**

Question	Answer	Marks																																												
2(c)	<table border="1" data-bbox="595 218 1677 842"> <thead> <tr> <th data-bbox="595 218 824 268">A</th> <th data-bbox="824 218 1048 268">B</th> <th data-bbox="1048 218 1272 268">C</th> <th data-bbox="1272 218 1677 268">D</th> </tr> <tr> <th data-bbox="595 268 824 387">Time / min</th> <th data-bbox="824 268 1048 387">volume of nitrogen, <math>V / \text{cm}^3</math></th> <th data-bbox="1048 268 1272 387"><math>V / V_{\text{FINAL}}</math></th> <th data-bbox="1272 268 1677 387"><math>[\text{C}_6\text{H}_5\text{N}_2^+\text{Cl}^-(\text{aq})] / \text{mol dm}^{-3}</math></th> </tr> </thead> <tbody> <tr><td>0.0</td><td>0</td><td>0.000</td><td>0.0150</td></tr> <tr><td>2.0</td><td>9</td><td>0.125</td><td>0.0131</td></tr> <tr><td>4.0</td><td>17</td><td>0.236</td><td>0.0115</td></tr> <tr><td>6.0</td><td>24</td><td>0.333</td><td>0.0100</td></tr> <tr><td>8.0</td><td>30</td><td>0.417</td><td>0.00875</td></tr> <tr><td>10.0</td><td>35</td><td>0.486</td><td>0.00771</td></tr> <tr><td>12.0</td><td>40</td><td>0.556</td><td>0.00666</td></tr> <tr><td>14.0</td><td>44</td><td>0.611</td><td>0.00584</td></tr> <tr><td>16.0</td><td>48</td><td>0.667</td><td>0.00500</td></tr> </tbody> </table>	A	B	C	D	Time / min	volume of nitrogen, $V / \text{cm}^3$	$V / V_{\text{FINAL}}$	$[\text{C}_6\text{H}_5\text{N}_2^+\text{Cl}^-(\text{aq})] / \text{mol dm}^{-3}$	0.0	0	0.000	0.0150	2.0	9	0.125	0.0131	4.0	17	0.236	0.0115	6.0	24	0.333	0.0100	8.0	30	0.417	0.00875	10.0	35	0.486	0.00771	12.0	40	0.556	0.00666	14.0	44	0.611	0.00584	16.0	48	0.667	0.00500	
A	B	C	D																																											
Time / min	volume of nitrogen, $V / \text{cm}^3$	$V / V_{\text{FINAL}}$	$[\text{C}_6\text{H}_5\text{N}_2^+\text{Cl}^-(\text{aq})] / \text{mol dm}^{-3}$																																											
0.0	0	0.000	0.0150																																											
2.0	9	0.125	0.0131																																											
4.0	17	0.236	0.0115																																											
6.0	24	0.333	0.0100																																											
8.0	30	0.417	0.00875																																											
10.0	35	0.486	0.00771																																											
12.0	40	0.556	0.00666																																											
14.0	44	0.611	0.00584																																											
16.0	48	0.667	0.00500																																											
	Column values for <b>D</b> correctly calculated	<b>1</b>																																												
	3 sf in <b>C</b> and <b>D</b>	<b>1</b>																																												
2(d)	Candidate's calculated points correctly plotted from table in <b>2(c)</b>	<b>1</b>																																												
	Smooth curve of best fit	<b>1</b>																																												
2(e)	Tangent drawn at time zero	<b>1</b>																																												
	2 sets of co-ordinates shown	<b>1</b>																																												
	calculation of gradient of tangent	<b>1</b>																																												
	$\text{mol dm}^{-3} \text{ minute(s)}^{-1}$	<b>1</b>																																												

**PUBLISHED**

Question	Answer					Marks															
2(f)	<table border="1"> <thead> <tr> <th data-bbox="638 217 875 268">concentration 1</th> <th data-bbox="875 217 1008 268">Time 1</th> <th data-bbox="1008 217 1254 268">concentration 2</th> <th data-bbox="1254 217 1456 268">time 2</th> <th data-bbox="1456 217 1635 268"><math>t_{1/2}</math></th> </tr> </thead> <tbody> <tr> <td data-bbox="638 268 875 319">(0.0120)</td> <td data-bbox="875 268 1008 319">3</td> <td data-bbox="1008 268 1254 319">(0.0060)</td> <td data-bbox="1254 268 1456 319">13.4</td> <td data-bbox="1456 268 1635 319">10.4</td> </tr> <tr> <td data-bbox="638 319 875 371">0.010</td> <td data-bbox="875 319 1008 371">6</td> <td data-bbox="1008 319 1254 371">0.005</td> <td data-bbox="1254 319 1456 371">16.0</td> <td data-bbox="1456 319 1635 371">10.0</td> </tr> </tbody> </table>					concentration 1	Time 1	concentration 2	time 2	$t_{1/2}$	(0.0120)	3	(0.0060)	13.4	10.4	0.010	6	0.005	16.0	10.0	
	concentration 1	Time 1	concentration 2	time 2	$t_{1/2}$																
	(0.0120)	3	(0.0060)	13.4	10.4																
	0.010	6	0.005	16.0	10.0																
Columns 1 and 3					<b>1</b>																
Columns 2 and 4					<b>1</b>																
Half-lives correctly calculated.					<b>1</b>																
2(g)	First order <b>AND</b> because half-lives are constant/equal					<b>1</b>															