



Mark Scheme (Results)

Summer 2018

Pearson Edexcel GCSE
Combined Science - Paper 3
Chemistry 1 (1SC0_1CH)

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2018

Publications Code 1SC0_1CH_1806_MS

All the material in this publication is copyright

© Pearson Education Ltd 2018

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Mark schemes have been developed so that the rubrics of each mark scheme reflects the characteristics of the skills within the AO being targeted and the requirements of the command word. So for example the command word 'Explain' requires an identification of a point and then reasoning/justification of the point.

Explain questions can be asked across all AOs. The distinction comes whether the identification is via a judgment made to reach a conclusion, or, making a point through application of knowledge to reason/justify the point made through application of understanding. It is the combination and linkage of the marking points that is needed to gain full marks.

When marking questions with a 'describe' or 'explain' command word, the detailed marking guidance below should be consulted to ensure consistency of marking.

Assessment Objective		Command Word	
Strand	Element	Describe	Explain
AO1*		An answer that combines the marking points to provide a logical description	An explanation that links identification of a point with reasoning/justification(s) as required
AO2		An answer that combines the marking points to provide a logical description, showing application of knowledge and understanding	An explanation that links identification of a point (by applying knowledge) with reasoning/justification (application of understanding)
AO3	1a and 1b	An answer that combines points of interpretation/evaluation to provide a logical description	
AO3	2a and 2b		An explanation that combines identification via a judgment to reach a conclusion via justification/reasoning
AO3	3a	An answer that combines the marking points to provide a logical description of the plan/method/experiment	
AO3	3b		An explanation that combines identifying an improvement of the experimental procedure with a linked justification/reasoning

*there will be situations where an AO1 question will include elements of recall of knowledge directly from the specification (up to a maximum of 15%). These will be identified by an asterisk in the mark scheme.

Question Number	Answer	Additional guidance	Mark
1(a)(i)	<p>A description including</p> <ul style="list-style-type: none"> • apply lighted splint (1) • (squeaky) pop (1) 	<p>allow flame / ignite gas / fire</p> <p>ignore 'squeaky pop test' / glowing splint</p> <p>second mark is dependent on first</p>	<p>(2)</p> <p>AO 2 2</p>

Question Number	Answer	Additional guidance	Mark
1(a)(ii)	<p>An explanation linking</p> <ul style="list-style-type: none"> • loss of electron(s) (1) • two electrons (1) 	<p>allow gains two electrons for 1 mark</p> <p>zero marks overall if sharing of electrons / gain or loss of protons / positive electrons</p> <p>marks can be awarded for suitably drawn diagram / half equation</p>	<p>(2)</p> <p>AO 1 1</p>

Question Number	Answer	Additional guidance	Mark
1(b)	<p>final answer of 94 (g dm⁻³) with or without working (2)</p> <p>OR</p> $\frac{23.5}{250} (1) (= 0.094)$ <p>0.094 x 1000 (1)</p> <p>OR</p> $\frac{250}{1000} (\text{dm}^3) (1) (= 0.25 (\text{dm}^3))$ <p>$\frac{23.5}{0.25} (1)$</p> <p>OR</p> $\frac{1000}{250} (1) = 4$ <p>4 x 23.5 (1)</p>	<p>allow ECF (error carried forward) throughout</p> <p>other final answers: 0.094 / 9.4 (1) 0.000094 or 9.4 x 10⁻⁵ (1)</p> <p>0.25 (dm³) (1)</p> <p>allow $\frac{250}{23.5} \times 1000$ or 10638(.3) (1)</p>	<p>(2)</p> <p>AO 2 1</p>

Question Number	Answer	Additional guidance	Mark
1(c)	<p>A description to include</p> <ul style="list-style-type: none"> • filter (1) <p>and two in a logical order from</p> <ul style="list-style-type: none"> • crystallisation (1) • heat solution (to concentrate) (1) • allow to cool (1) • dry crystals between filter papers (1) 	<p>if filtration not first stage, ignore it and give maximum 2 marks</p> <p>allow description of filtration ignore filtration to obtain nickel sulfate (crystals)</p> <p>allow 'leave until water evaporates' / use of water bath / evaporate {water/the solution}</p> <p>allow leave {until crystals form / for a few hours / in a warm place / on a window sill}</p> <p>allow 'dry crystals in (warm) oven'</p> <p>if alternative methods of making nickel sulfate solution described, max 1 mark from last four marking points</p>	<p>(3)</p> <p>AO 2 2</p>

Total for question 1 = 9 marks

Question Number	Answer	Mark
2(a)(i)	<p data-bbox="288 197 671 226">C iron oxide is reduced</p> <p data-bbox="288 264 759 293">The only correct answer is C</p> <p data-bbox="288 331 970 360"><i>A is not correct because carbon gains oxygen</i></p> <p data-bbox="288 398 1102 427"><i>B is not correct because it is not an acid-base reaction</i></p> <p data-bbox="288 465 1015 495"><i>D is not correct because iron oxide loses oxygen</i></p>	<p data-bbox="1374 197 1430 226">(1)</p> <p data-bbox="1374 264 1477 293">AO 1 1</p>

Question Number	Answer	Additional guidance	Mark
2(a)(ii)	<p>final answer of 168 (tonnes) with or without working (3)</p> <p>OR</p> <p>relative formula mass $\text{Fe}_2\text{O}_3 = 2 \times 56 + 3 \times 16 (= 160)$ (1)</p> <p>160 tonnes Fe_2O_3 produces $\{2 \times 56 / 112\}$ tonnes Fe (1)</p> <p>240 tonnes Fe_2O_3 produces $\frac{2 \times 56}{160} \times 240$ (1) = 168 (tonnes)</p> <p>OR</p> <p>relative formula mass $\text{Fe}_2\text{O}_3 = 2 \times 56 + 3 \times 16 (= 160)$ (1)</p> <p>$\frac{240}{160}$ (1) = 1.5</p> <p>1.5 x 112 (1) = 168 (tonnes)</p> <p>OR</p> <p>relative formula mass $\text{Fe}_2\text{O}_3 = 2 \times 56 + 3 \times 16 (= 160)$ (1)</p> <p>$\frac{112}{160}$ (1) = 0.7</p> <p>0.7 x 240 (1) = 168 (tonnes)</p>	<p>allow ECF throughout</p> <p>$M_r [\text{Fe}_2\text{O}_3] = 160$ seen without working (1)</p> <p>allow 320 tonnes : 224 tonnes (1)</p> <p>final answer 84 (tonnes) with or without working (2)</p> <p>Note : final answer 1.5 scores 2 overall</p>	<p>(3)</p> <p>AO 2 1</p>

Question Number	Answer	Additional guidance	Mark
2(b)	<p>An explanation linking the following points</p> <ul style="list-style-type: none"> aluminium is high in reactivity / aluminium oxide is (very) stable (1) aluminium (oxide) cannot be reduced by carbon (1) 	<p>allow carbon is less reactive than aluminium / ORA / aluminium is very reactive ignore 'aluminium is more reactive' (alone)</p> <p>allow carbon cannot displace aluminium / aluminium oxide does not react with carbon</p> <p>ignore aluminium extracted by electrolysis</p>	<p>(2)</p> <p>AO 1 1</p>

Question Number	Answer	Mark
2(c)	electrolysis	<p>(1)</p> <p>AO 3 2a</p>

Question Number	Answer	Additional guidance	Mark
2(d)	<p>A description to include</p> <ul style="list-style-type: none"> plants absorb {copper/metal} (ions) from the {soil/ores} / plants concentrate copper ions (1) plants (harvested and) burned (to leave copper/metal compound) (1) 	<p>ignore plants absorb copper from solid metal ignore copper {atoms/metal/compounds}</p> <p>ignore plants heated</p> <p>mark independently</p>	<p>(2)</p> <p>AO 1 1</p>

Total for question 2 = 9 marks

Question Number	Answer	Mark
3(a)	<p>C 30 2403</p> <p>The only correct answer is C</p> <p><i>A is not correct because it will be a solid above 80 °C</i></p> <p><i>B is not correct because it will be a liquid at 20 °C and gas at 80 °C</i></p> <p><i>D is not correct because it will be a liquid at 20 °C and gas at 80 °C</i></p>	<p>(1)</p> <p>AO 1 1</p>

Question Number	Answer	Additional guidance	Mark
3(b)(i)	<p>An explanation linking</p> <ul style="list-style-type: none"> water {boils / evaporates} (to form steam / water vapour / leaving salt behind) (1) (steam / water vapour) condenses (to form pure water) (1) <p>allow alternative wording for evaporate and condense</p>	<p>ignore sea water evaporates</p> <p>sea water evaporates and condenses scores 1 overall</p> <p>mark independently</p>	<p>(2)</p> <p>AO 1 1</p>

Question Number	Answer	Additional guidance	Mark
3(b)(ii)	<p>An explanation linking</p> <ul style="list-style-type: none"> use a (Liebig) condenser / surround test tube with (beaker of) {iced/cold} water / wrap delivery tube with cold cloth (1) to increase effectiveness of cooling / amount of condensation / remove the heat energy more effectively / ensure all the water (vapour) condenses (1) 	<p>ignore anti bumping granules / fractionating column</p> <p>allow alternative suitably described methods / prevent water vapour escaping / cools water vapour faster</p> <p>ignore sea water vapours</p> <p>a closed system scores 0 overall</p> <p>mark independently</p>	<p>(2)</p> <p>AO 3 3b</p>

Question Number	Answer	Additional guidance	Mark
3(c)	<p>An explanation linking</p> <p>from B to C / at B: graph flat because</p> <ul style="list-style-type: none"> • particles in solid use energy to {break out of lattice / break (intermolecular) bonds (between particles) / particles becoming randomly arranged / turn solid to liquid} (1) <p>and any three from</p> <p>from A to B / at A: graph rises because</p> <ul style="list-style-type: none"> • particles in solid in a lattice / fixed (mean) positions (1) • vibrate more (rapidly) (as temperature increases) (1) <p>from C to D / at C: graph rises because</p> <ul style="list-style-type: none"> • particles in liquid move past one another / randomly (1) • particles move more (rapidly) (as temperature increases) (1) 	<p>may be shown as a diagram / on graph</p> <p>may be shown as a diagram / on graph ignore references to gas / evaporation / boil</p>	<p>(4)</p> <p>AO 3 2a AO 3 2b</p>

Total for question 3 = 9 marks

Question Number	Answer	Mark
4(a)(i)	<p>C chlorine zinc</p> <p>The only correct answer is C</p> <p><i>A is not correct because oxygen cannot be produced by the electrolysis of this molten salt</i></p> <p><i>B is not correct because hydrogen cannot be produced by the electrolysis of this molten salt</i></p> <p><i>D is not correct because hydrogen and oxygen cannot be produced by the electrolysis of this molten salt</i></p>	<p>(1)</p> <p>AO 2 1</p>

Question Number	Answer	Mark
4(a)(ii)	<p>D it contains ions that can move</p> <p>The only correct answer is D</p> <p><i>A is not correct because molten zinc chloride does not contain molecules</i></p> <p><i>B is not correct because molten zinc chloride does not have a giant structure</i></p> <p><i>C is not correct because delocalised electrons are not present</i></p>	<p>(1)</p> <p>AO 1 1</p>

Question Number	Answer	Additional guidance	Mark
4(b)(i)	<p>A diagram of a workable apparatus showing a complete circuit including</p> <ul style="list-style-type: none"> • electrodes labelled in (copper sulfate) solution (1) • {power supply / power pack / battery} connected (1) 	<p>max 1 if circuit not complete</p> <p>allow labelling as 'electrodes' or 'anode' and 'cathode' or 'copper'</p> <p>ignore 'connected to mains'</p> <p>allow symbol for cell/battery even if wrong way round</p>	<p>(2)</p> <p>AO 1 2</p>


Question Number	Answer	Additional guidance	Mark
4(b)(ii)	<p>An explanation linking the following point to a maximum of four</p> <ul style="list-style-type: none"> anode lost copper and cathode gained copper / reaction at cathode is reverse of reaction at anode / copper ions move into solution at anode AND copper ions move out of solution at cathode (1) <p>and any three from</p> <ul style="list-style-type: none"> at anode copper atoms become copper ions (1) and lose two electrons (1) OR (at anode) $\text{Cu} \rightarrow \text{Cu}^{2+} + 2\text{e}^-$ (2) at cathode copper ions become copper atoms (1) and gain two electrons (1) OR (at cathode) $\text{Cu}^{2+} + 2\text{e}^- \rightarrow \text{Cu}$ (2) 	<p>ignore references to zinc, chlorine and zinc chloride</p> <p>allow copper atoms are oxidised (1)</p> <p>marking points independently</p> <p>allow copper ions are reduced (1)</p> <p>marking points independently</p> <p>penalise wrong use of atom / ion once only</p> <p>penalise wrong use of reduced / oxidised once only</p>	<p>(4)</p> <p>AO 2 1</p>

Question Number	Answer	Additional guidance	Mark
4(c)	<p>$2\text{H}^+ + 2\text{e}^{(-)} \rightarrow \text{H}_2$ / $2\text{H}^+ \rightarrow \text{H}_2 - 2\text{e}^{(-)}$ (2)</p> <p>species in correct place as shown above (1)</p> <p>balancing of correct species in correct place (1)</p>	<p>allow use of = or \rightleftharpoons in place of \rightarrow</p> <p>allow multiples</p> <p>reject h_2 / h_2 / H_2 / H^2</p>	<p>(2)</p> <p>AO 1 1</p>

Total for question 4 = 10 marks

Question Number	Answer	Mark
5(a)(i)	<p>B -78 -33 does not conduct</p> <p>The only correct answer is B</p> <p><i>A is not correct because simple molecular, covalent substances do not have high mpt and bpt</i></p> <p><i>C is not correct because ammonia is a gas at room temperature and does not conduct</i></p> <p><i>D is not correct because simple molecular, covalent substances do not have these properties</i></p>	<p>(1)</p> <p>AO 2 1</p>

Question Number	Answer	Additional guidance	Mark
5(a)(ii)	<p>$N_2 + 3H_2 \rightarrow 2NH_3$ (2)</p> <p>left hand side formulae (1)</p> <p>balancing of correct formulae (1)</p>	<p>accept multiples</p> <p>allow = or \rightleftharpoons in place of \rightarrow</p> <p>ignore state symbols even if incorrect</p> <p>do not allow N2, n2, etc</p>	<p>(2)</p> <p>AO 2 1</p>


Question Number	Answer	Additional guidance	Mark
5(b)	 <p>(2)</p>	<p>double bond (1)</p> <p>rest of molecule (1)</p> <p>(dependent on correct double bond)</p> <p>ignore atomic symbol</p> <p>allow all x or •</p> <p>ignore inner shells of electrons even if incorrect</p>	<p>(2)</p> <p>AO 1 1</p>

Question Number	Indicative content	Mark
5(c)	<p>Answers will be credited according to candidate's deployment of knowledge and understanding of the material in relation to the qualities and skills outlines in the generic mark scheme.</p> <p>The indicative content below is not prescriptive and candidates are not required to include all the material which is indicated as relevant. Additional content included in the response must be scientific and relevant.</p> <ul style="list-style-type: none"> • in all structures the carbon atoms bonded by single covalent bonds • shared pair of electrons • strong bonds • in diamond each carbon atom joined to four others • diamond has a giant covalent {structure/lattice} • graphene has a giant covalent {structure/lattice} • fullerene has a molecular structure • in graphene and fullerene each carbon atom joined to three others • in diamond and graphene many bonds need to be broken to melt • need lots of energy • therefore very high melting / sublimation points • in fullerene weak forces between molecules • less energy needed to separate molecules • fullerene has the lowest melting / sublimation point • because diamond and graphene have lots of strong covalent bonds so both are very strong materials • because weak forces between fullerene molecules so its strength is very low • in diamond there are no free electrons • so diamond does not conduct • in graphene and fullerene each carbon atom has one free electron • hence delocalised electrons • graphene conducts electricity • fullerene only conducts electricity across the surface of the molecule • no/little movement of electrons between molecules • so fullerene is poor conductor of electricity (/ semi conductor) 	(6) AO 1 1
Level	Mark	Descriptor
	0	No rewardable material.
Level 1	1–2	<ul style="list-style-type: none"> • Demonstrates elements of chemical understanding, some of which is inaccurate. Understanding of scientific ideas, enquiry, techniques and procedures lacks detail. (AO1) • Presents an explanation with some structure and coherence. (AO1)
Level 2	3–4	<ul style="list-style-type: none"> • Demonstrates chemical understanding, which is mostly relevant but may include some inaccuracies. Understanding of scientific ideas, enquiry, techniques and procedures is not fully detailed and fully devolved. (AO1)

		<ul style="list-style-type: none">• Presents an explanation that has a structure which is mostly clear, coherent and logical. (AO1)
Level 3	5–6	<ul style="list-style-type: none">• Demonstrates accurate and relevant chemical understanding throughout. Understanding of the scientific ideas, enquiry, techniques and procedures is detailed and fully devolved. (AO1)• Presents an explanation that has a well-developed structure which is clear, coherent and logical. (AO1)

Total for question 5 = 11 marks

Question Number	Answer	Additional guidance	Mark
6(a)(i)	P R Q S (2)	two in correct order (1)	(2) AO 3 2a AO 3 2b

Question Number	Answer	Additional guidance	Mark
6(a)(ii)	<p>A workable diagram showing a method to measure the volume of the gas</p> <ul style="list-style-type: none"> delivery tube between test-tube and (1) gas syringe / (graduated tube / inverted burette / measuring cylinder) over water bath (1) 	<p>if diagram is not workable (e.g. no bung at top of test tube), max 1 mark</p> <p>allow connection shown as</p>  <p>if collection vessel not labelled, graduations must be shown for the second mark</p>	(2) AO 3 3a AO 3 3b

Question Number	Answer	Additional guidance	Mark
6(b)	<p>iron $\frac{10.00}{56} = 0.179 / 0.18 / 0.2$ and</p> <p>copper $\frac{11.34}{63.5} = 0.179 / 0.18 / 0.2$ (1)</p> <p>(ratio 1:1) so reaction A (1)</p>	<p>allow max 1 mark for</p> <p>Fe : $\frac{56}{10.00} = 5.6$</p> <p>Cu : $\frac{63.5}{11.34} = 5.6$ so reaction A</p> <p>other methods of calculation include</p> <p>10.00 g Fe forms $\frac{10.00}{56} \times 63.5$ (1) g copper</p> <p>$= 11.34$ g</p> <p>copper so reaction A (1)</p> <p>second mark dependent on first</p>	(2) AO 3 2a AO 3 2b

Question Number	Answer	Additional guidance	Mark
6(c)	$2\text{Al} + 6\text{H}^+ \rightarrow 2\text{Al}^{3+} + 3\text{H}_2$ (2)	Al and H ₂ (1) balancing of correct species (1) allow multiples	(2) AO 2 1

Question Number	Answer	Additional guidance	Mark
6(d)	pH {increases / goes up} by <u>one</u> / moves <u>1</u> closer to neutral	ignore {increases / goes up} alone	(1) AO 1 1

Question Number	Answer	Additional guidance	Mark
6(e)	<p>1 mol of hydrogen atoms = a mass of 1.00 g = 6.02×10^{23} atoms</p> <p>6.02×10^{23} H atoms has mass = 1.00 g (1)</p> <p>mass of 1 H atom = <u>1.00</u> (1)</p> $= \frac{6.02 \times 10^{23}}{6.02 \times 10^{23}} = 1.66 \times 10^{-24}$ <p>(g) (1)</p>	<p>correct answer alone (3)</p> <p>if $1 \times 6.02 \times 10^{23}$ is followed by atoms or particles, then award 1st marking point</p> <p>on answer line 3.32×10^{-24} (g) (2)</p> <p>ignore sig figs except for one</p>	<p>(3)</p> <p>AO 2 1</p>

Total for question 6 = 12 marks

