



Mark Scheme (Results)

January 2021

Pearson Edexcel International Advanced Level
In Chemistry (WCH14)
Paper 1: Rates, Equilibria and Further Organic
Chemistry

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

January 2021

Publications Code WCH14_01_2101_MS

All the material in this publication is copyright

© Pearson Education Ltd 2021

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Using the mark scheme

Examiners should look for qualities to reward rather than faults to penalise. This does NOT mean giving credit for incorrect or inadequate answers, but it does mean allowing candidates to be rewarded for answers showing correct application of principles and knowledge. Examiners should therefore read carefully and consider every response: even if it is not what is expected it may be worthy of credit.

The mark scheme gives examiners:

- an idea of the types of response expected
- how individual marks are to be awarded
- the total mark for each question
- examples of responses that should NOT receive credit.

/ means that the responses are alternatives and either answer should receive full credit. () means that a phrase/word is not essential for the award of the mark, but helps the examiner to get the sense of the expected answer.

Phrases/words in **bold** indicate that the meaning of the phrase or the actual word is **essential** to the answer. ecf/TE/cq (error carried forward) means that a wrong answer given in an earlier part of a question is used correctly in answer to a later part of the same question.

Candidates must make their meaning clear to the examiner to gain the mark. Make sure that the answer makes sense. Do not give credit for correct words/phrases which are put together in a meaningless manner. Answers must be in the correct context.

Quality of Written Communication

Questions which involve the writing of continuous prose will expect candidates to:

- write legibly, with accurate use of spelling, grammar and punctuation in order to make the meaning clear
- select and use a form and style of writing appropriate to purpose and to complex subject matter
- organise information clearly and coherently, using specialist vocabulary when appropriate.

Full marks will be awarded if the candidate has demonstrated the above abilities.

Questions where QWC is likely to be particularly important are indicated (QWC) in the mark scheme, but this does not preclude others.

Section A

Question Number	Answer	Mark
1	<p>The only correct answer is A (carbon dioxide, CO₂)</p> <p><i>B is not correct because copper is a solid at 298 K and 1 atm pressure so has the lowest entropy</i></p> <p><i>C is not correct because ethanol is a liquid at 298 K and 1 atm pressure and has a lower entropy than a gas</i></p> <p><i>D is not correct because hydrogen is also a gas at 298 K and 1 atm pressure, but its molecules are smaller than carbon dioxide molecules</i></p>	(1)

Question Number	Answer	Mark
2	<p>The only correct answer is B (reactions P and Q only)</p> <p><i>A is not correct because both reactions P and Q have a positive value for ΔS_{total}</i></p> <p><i>C is not correct because reaction R has a negative value for ΔS_{total} so is not feasible</i></p> <p><i>D is not correct because both reactions R and S have a negative value for ΔS_{total} so are not feasible</i></p>	(1)

Question Number	Answer	Mark
3	<p>The only correct answer is A ($\frac{1}{2}\text{Br}_2(\text{l}) \rightarrow \text{Br}(\text{g})$)</p> <p><i>B is not correct because the standard enthalpy change of atomisation refers to the formation of 1 mol of atoms</i></p> <p><i>C is not correct because bromine's standard state is as a liquid</i></p> <p><i>D is not correct because bromine exists as diatomic molecules in the liquid state in its standard state and only 1 mol of atoms should be formed</i></p>	(1)

Question Number	Answer	Mark
4(a)	<p>The only correct answer is C (potassium bromide)</p> <p><i>A is not correct because the least exothermic lattice energy is between the largest ions with the smallest charge and calcium ions are smaller and have a higher charge than potassium ions</i></p> <p><i>B is not correct because the least exothermic lattice energy is between the largest ions with the smallest charge and magnesium ions are smaller and have a higher charge than potassium ions</i></p> <p><i>D is not correct because sodium ions are smaller than potassium ions</i></p>	(1)

Question Number	Answer	Mark
4(b)	<p>The only correct answer is B (magnesium chloride)</p> <p><i>A is not correct because Ca^{2+} ions are larger than magnesium ions so will polarise less</i></p> <p><i>C is not correct because K^+ ions are larger than magnesium ions and have a lower charge so will polarise less</i></p> <p><i>D is not correct because Na^+ ions are larger than magnesium ions and have a lower charge so will polarise less</i></p>	(1)

Question Number	Answer	Mark
5	<p>The only correct answer is C (the total entropy when KCl dissolves is positive)</p> <p><i>A is not correct because the enthalpy change of hydration for all ions and lattice energy for all ionic compounds are exothermic so this does not explain why KCl is soluble</i></p> <p><i>B is not correct because the enthalpy change of hydration for all ions and lattice energy for all ionic compounds are exothermic</i></p> <p><i>D is not correct because the total entropy must be positive for a spontaneous reaction</i></p>	(1)

Question Number	Answer	Mark
6	<p>The only correct answer is A (3.61×10^{-5})</p> <p><i>B is not correct because R and ΔS_{total} are the wrong way up</i></p> <p><i>C is not correct because the temperature should not be included</i></p> <p><i>D is not correct because the negative sign for ΔS_{total} has been omitted</i></p>	(1)

Question Number	Answer	Mark
7	<p>The only correct answer is A $(\text{CH}_3\text{COCH}_3 + \text{H}^+ \rightleftharpoons \text{CH}_3\text{C}(\text{O}^+\text{H})\text{CH}_3$ fast, $\text{CH}_3\text{C}(\text{O}^+\text{H})\text{CH}_3 \rightarrow \text{CH}_3\text{C}(\text{OH})=\text{CH}_2 + \text{H}^+$ slow, $\text{CH}_3\text{C}(\text{OH})=\text{CH}_2 + \text{I}_2 \rightarrow \text{CH}_3\text{COCH}_2\text{I} + \text{HI}$ fast)</p> <p><i>B is not correct because the steps up to and including the slow step must include CH_3COCH_3 and H^+ ions and I_2 must only be involved in a fast step</i></p> <p><i>C is not correct because the steps up to and including the slow step must include CH_3COCH_3 and H^+ ions and I_2 must only be involved in a fast step</i></p> <p><i>D is not correct because the steps up to and including the slow step must include CH_3COCH_3 and H^+ ions and I_2 must only be involved in a fast step</i></p>	(1)

Question Number	Answer	Mark
8	<p>The only correct answer is C (1.0×10^{-3})</p> <p><i>A is not correct because the initial rate and the rate constant have been mixed up</i></p> <p><i>B is not correct because $[A]$ has not been squared</i></p> <p><i>D is not correct because $[B]$ has been included</i></p>	(1)

Question Number	Answer	Mark
9(a)	<p>The only correct answer is B (1.95)</p> <p><i>A is not correct because pK_a has not been converted to K_a</i></p> <p><i>C is not correct because this is the pH of $0.100 \text{ mol dm}^{-3}$ ethanoic acid</i></p> <p><i>D is not correct because the square root of $K_a \times [\text{CH}_2\text{ClCOOH}]$ has not been used to calculate $[\text{H}^+]$</i></p>	(1)

Question Number	Answer	Mark
9(b)	<p>The only correct answer is C (Acid: CH_2ClCOOH, Conjugate base: $\text{CH}_2\text{ClCOO}^-$)</p> <p><i>A is not correct because ethanoic acid has a higher pK_a than chloroethanoic acid so acts as a base in this reaction</i></p> <p><i>B is not correct because ethanoic acid has a higher pK_a than chloroethanoic acid so acts as a base in this reaction</i></p> <p><i>D is not correct because chloroethanoic acid loses a proton when it acts as an acid</i></p>	(1)


Question Number	Answer	Mark
10(a)	<p>The only correct answer is B (region U)</p> <p><i>A is not correct because in region T there is only aqueous ammonia at the start of the titration</i></p> <p><i>C is not correct because in region V, the vertical part of the graph, represents the end-point of the titration</i></p> <p><i>D is not correct because in region W all the aqueous ammonia has been neutralised</i></p>	(1)

Question Number	Answer	Mark
10(b)	<p>The only correct answer is A (methyl red)</p> <p><i>B is not correct because phenol red has a pH range of 6.8 to 8.4 and 8.4 is not in the vertical region</i></p> <p><i>C is not correct because phenolphthalein has a pH range of 8.2 to 10.0 and this is not in the vertical region</i></p> <p><i>D is not correct because thymol blue has a pH range of 1.2 to 2.8 and this is not in the vertical region</i></p>	(1)

Question Number	Answer	Mark
10(c)	<p>The only correct answer is B (5.8)</p> <p><i>A is not correct because this is the approximate pH when excess hydrochloric acid has been added</i></p> <p><i>C is not correct because this is the approximate pH near the start of the end point when there is still excess aqueous ammonia</i></p> <p><i>D is not correct because this is the approximate pH of aqueous ammonia</i></p>	(1)

Question Number	Answer	Mark
11	<p>The only correct answer is B (9.77×10^{-2} (mol dm⁻³))</p> <p><i>A is not correct because the two volumes have been reversed</i></p> <p><i>C is not correct because a mole ratio of 1:1 has been used instead of 1 mol of acid : 2 mol NaOH</i></p> <p><i>D is not correct because a mole ratio of 2 mol of acid : 1 mol NaOH has been used</i></p>	(1)

Question Number	Answer	Mark
12	<p>The only correct answer is C (tertiary only)</p> <p><i>A is not correct because primary bromoalkanes react by an S_N2 mechanism and the product would be optically active</i></p> <p><i>B is not correct because secondary bromoalkanes react by an S_N1 or an S_N2 mechanism and the product could be optically active</i></p> <p><i>D is not correct because primary and secondary bromoalkanes react by S_N1 and S_N2 mechanisms and the products could be optically active</i></p>	(1)

Question Number	Answer	Mark
13	<p>The only correct answer is D</p>  <p><i>A is not correct because this is an aldehyde and it would give a silver mirror with Tollens' reagent</i></p> <p><i>B is not correct because this is an aldehyde and it would give a silver mirror with Tollens' reagent</i></p> <p><i>C is not correct because this is an aldehyde and it would give a silver mirror with Tollens' reagent</i></p>	(1)

Question Number	Answer	Mark
14	<p>The only correct answer is C (CH₃COONa and CH₃CH₂CH₂OH)</p> <p><i>A is not correct because ethanoic acid reacts with NaOH to form the sodium salt</i></p> <p><i>B is not correct because ethanoic acid reacts with NaOH to form the sodium salt and propan-1-ol does not react with NaOH</i></p> <p><i>D is not correct because propan-1-ol does not react with NaOH</i></p>	(1)

Question Number	Answer	Mark
15	<p>The only correct answer is C (90.5 %)</p> <p><i>A is not correct because the molar masses have been used for the incorrect substances and the amount of ethanoic acid should be the numerator of the fraction</i></p> <p><i>B is not correct because the masses have not been converted into moles and the amount of ethanoic acid should be the numerator of the fraction</i></p> <p><i>D is not correct because the masses have not been converted into moles</i></p>	(1)

Question Number	Answer	Mark
16	<p>The only correct answer is D (weak attraction to stationary phase, strong attraction to mobile phase)</p> <p><i>A is not correct because if there was a strong attraction to the stationary phase the component would not move very far and would have a low R_f value</i></p> <p><i>B is not correct because if there was a weak attraction to the mobile phase the component would not move very far and would have a low R_f value</i></p> <p><i>C is not correct because if there was a weak attraction to the mobile phase the component would not move very far and would have a low R_f value</i></p>	(1)



(Total for Section A = 20 marks)

Section B

Question Number	Answer	Additional Guidance	Mark
17(a)(i)	<p>An answer that makes reference to the following point:</p> <ul style="list-style-type: none"> it / lactic acid is non-superimposable on its mirror image 	<p>Allow there are four different atoms / groups attached to a carbon (atom)</p> <p>Allow it is chiral / has a chiral centre / has a chiral carbon (atom) / has an asymmetric carbon (atom)</p> <p>Ignore rotates the plane of plane-polarised light</p> <p>Do not award four different molecules attached to a carbon (atom)</p>	(1)

Question Number	Answer	Additional Guidance	Mark
17(a)(ii)	<p>An answer that makes reference to the following point:</p> <ul style="list-style-type: none"> it is a racemic mixture or contains equal amounts of the two enantiomers / (optical) isomers 	<p>Allow rotations caused by both enantiomers / isomers cancel</p> <p>Ignore just contains two enantiomers / isomers</p> <p>Do not award plane-polarised light cannot pass through the solution</p>	(1)

Question Number	Answer	Additional Guidance	Mark
17(a)(iii)	<ul style="list-style-type: none"> $\text{CH}_2=\text{CHCOOH}$ 	<p>Allow any combination of structural or displayed formulae or skeletal formulae / $\text{COOHCH}=\text{CH}_2$</p>	(1)


Question Number	Answer	Additional Guidance	Mark
17(b)	<ul style="list-style-type: none"> • X: butan-1-ol or  or CH₃CH₂CH₂CH₂OH / CH₃(CH₂)₂CH₂OH (1) • Y: phosphorus(V) chloride / phosphorus pentachloride / PCl₅ (1) • Z: <i>N</i>-methylbutanamide or  or CH₃CH₂CH₂CONHCH₃ / CH₃(CH₂)₂CONHCH₃ (1) 	<p>Mark independently</p> <p>If names and formulae are given, both must be correct but penalise missing H from carbon chain displayed formulae once only</p> <p>Allow any combination of skeletal, structural or displayed formulae</p> <p>Ignore molecular formulae for X, Y and Z</p> <p>Ignore butanol / C₄H₉OH Do not award CH₃CH₂CH₂CH₃O Do not award butanal</p> <p>(1)</p> <p>Allow phosphorus(III) chloride / PCl₃ / thionyl chloride / SOCl₂ Do not award hydrochloric acid / HCl</p> <p>(1)</p> <p>Ignore methylbutanamide / butanamide in addition to a correct structure</p> <p>Allow NH in skeletal formula</p> <p>Do not award CH₃CH₂CH₂COHNCH₃ / CH₃CH₂CH₂COCH₃NH</p>	(3)

Question Number	Answer	Additional Guidance	Mark
17(c)	<p>• Monomer 1</p> $ \begin{array}{c} \text{H} \quad \text{H} \\ \quad \\ \text{H}-\text{O}-\text{C}-\text{C}-\text{O}-\text{H} \\ \quad \\ \text{H} \quad \text{H} \end{array} $ <p>(1)</p> <p>• Monomer 2</p> $ \begin{array}{c} \text{O} \quad \text{H} \quad \text{O} \\ \quad \quad \\ \text{H}-\text{O}-\text{C}-\text{C}=\text{C}-\text{C}-\text{O}-\text{H} \\ \quad \quad \\ \quad \quad \text{H} \end{array} $ <p>or</p> $ \begin{array}{c} \text{O} \quad \text{H} \quad \text{O} \\ \quad \quad \\ \text{Cl}-\text{C}-\text{C}=\text{C}-\text{C}-\text{Cl} \\ \quad \quad \\ \quad \quad \text{H} \end{array} $ <p>(1)</p>	<p>Allow monomers in either order Allow any combination of structural or displayed formulae / skeletal formulae Allow OH</p> <p>Ignore bond lengths and bond angles</p> <p>Penalise OH-C on left of molecules once only Penalise missing H from carbon chain displayed formulae once only</p> <p>Accept cis isomers</p>	(2)

Question Number	Answer	Additional Guidance	Mark
17(d)(i)	<ul style="list-style-type: none"> $C_5H_{10}O_2$ 	Allow symbols in any order Ignore any working Ignore + charge	(1)

Question Number	Answer	Additional Guidance	Mark
17(d)(ii)	<ul style="list-style-type: none"> E is not a carboxylic acid or does not contain COOH (group) 	If name and formula are given, both must be correct Allow E is not an acid Do not award additional functional groups	(1)

Question Number	Answer	Additional Guidance	Mark
17(d)(iii)	<ul style="list-style-type: none"> E is an ester 	Ignore saturated / -COO- / C=O Do not award additional functional groups	(1)

Question Number	Answer	Additional Guidance	Mark
17 (d)(iv)	<ul style="list-style-type: none"> • structure of E • 2 or 3 proton environments correct • 4th proton environment correct 	<p>Example of structure:</p>  <p>(1)</p> <p>(1)</p> <p>(1)</p> <p>Protons can be circled and labelled or just labelled Allow labels using data from the table</p> <p>Only 1 proton from each group needs to be labelled</p> <p>Allow whole groups to be labelled, including the carbon atom</p>	(3)

(Total for Question 17 = 14 marks)

Question Number	Answer	Additional Guidance	Mark
18(a)(i)	<ul style="list-style-type: none"> no effect / none / nothing / no change 	Ignore references to rate	(1)

Question Number	Answer	Additional Guidance	Mark
18(a)(ii)	<p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> (equilibrium) yield (of sulfur trioxide / SO_3 / product) decreases (1) the equilibrium constant / K_p / K_c / K decreases (as temperature increases) and because the (forward / right) reaction is exothermic / releases heat (energy) / ΔH is negative (1) 	<p>Allow less sulfur trioxide / SO_3 / product forms Ignore equilibrium position shifts to the left Ignore more reactants formed</p> <p>Allow the equilibrium constant / K_p / K_c / K decreases (as temperature increases) and because the reverse / backward / left reaction is endothermic / absorbs heat (energy)</p> <p>Allow K decreases because $\Delta S_{\text{surroundings}} / \Delta S_{\text{total}}$ decreases / becomes less positive (as temperature increases and assuming ΔS_{system} is constant)</p> <p>Ignore reference to rate</p>	(2)

Question Number	Answer	Additional Guidance	Mark
18(a)(iii)	<ul style="list-style-type: none"> expression for K_p 	<p>Example of expression for K_p:</p> $K_p = \frac{p(\text{SO}_3(\text{g}))^2}{p(\text{SO}_2(\text{g}))^2 \times p(\text{O}_2(\text{g}))}$ <p>Allow $P / PP / pp$ etc for partial pressure and this can be inside the brackets Allow e.g. $p^2\text{SO}_3$</p> <p>Ignore missing (g) / brackets around formulae</p> <p>Do not award square brackets</p>	(1)

Question Number	Answer	Additional Guidance	Mark																				
18(a)(iv)	<ul style="list-style-type: none"> • calculation of eqm moles (1) • calculation or expressions for 3 partial pressures (1) • substitution of values into K_p expression (1) • calculation of K_p and answer to 2 / 3 SF and units (1) 	<p>Example of calculation:</p> <table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <thead> <tr> <th style="width: 25%;"></th> <th style="width: 25%;">SO₂</th> <th style="width: 25%;">O₂</th> <th style="width: 25%;">SO₃</th> </tr> </thead> <tbody> <tr> <td>Initial mol</td> <td>2.00</td> <td>1.00</td> <td>-</td> </tr> <tr> <td>Eqm mol</td> <td>2.00 – 1.60 = 0.40</td> <td>1.00 – 0.80 = 0.20</td> <td>1.60</td> </tr> <tr> <td>Total mol at eqm</td> <td colspan="3">0.40 + 0.20 + 1.60 = 2.20</td> </tr> <tr> <td>Partial pressure /atm</td> <td>$\frac{0.40 \times 5.00}{2.20}$ = 0.90909</td> <td>$\frac{0.20 \times 5.00}{2.20}$ = 0.45455</td> <td>$\frac{1.60 \times 5.00}{2.20}$ = 3.6364</td> </tr> </tbody> </table> <p>TE for partial pressures on eqm moles</p> $K_p = \frac{3.6364^2}{0.90909^2 \times 0.45455}$ $= 35.2 / 35 \text{ atm}^{-1}$ <p>TE on expression for K_p in (a)(iii)</p> <p>Allow answer from previous correct rounding to 2 or more SF e.g. 0.91, 0.45 and 3.6 gives 34.8 Penalise incorrect rounding once only e.g. 0.909 to 0.9</p> <p>Allow fractions in working but not in final answer</p> <p>Allow atm⁻¹ for units but do not allow any other units Allow correct units written in (a)(iii) if not written here</p> <p>Correct answer to 2 or 3 SF with units and no working scores (4)</p>		SO ₂	O ₂	SO ₃	Initial mol	2.00	1.00	-	Eqm mol	2.00 – 1.60 = 0.40	1.00 – 0.80 = 0.20	1.60	Total mol at eqm	0.40 + 0.20 + 1.60 = 2.20			Partial pressure /atm	$\frac{0.40 \times 5.00}{2.20}$ = 0.90909	$\frac{0.20 \times 5.00}{2.20}$ = 0.45455	$\frac{1.60 \times 5.00}{2.20}$ = 3.6364	(4)
	SO ₂	O ₂	SO ₃																				
Initial mol	2.00	1.00	-																				
Eqm mol	2.00 – 1.60 = 0.40	1.00 – 0.80 = 0.20	1.60																				
Total mol at eqm	0.40 + 0.20 + 1.60 = 2.20																						
Partial pressure /atm	$\frac{0.40 \times 5.00}{2.20}$ = 0.90909	$\frac{0.20 \times 5.00}{2.20}$ = 0.45455	$\frac{1.60 \times 5.00}{2.20}$ = 3.6364																				

Question Number	Answer	Additional Guidance	Mark
18(b)(i)	<ul style="list-style-type: none"> • calculation of mass of H₂SO₄ in 1 dm³ • calculation of concentration of acid 	<p>Example of calculation: mass of H₂SO₄ in 1 dm³ concentrated acid = 0.985 x 1800 = 1773 (g)</p> <p>(1) concentration of acid = $\frac{1773}{(2 \times 1.0) + 32.1 + (4 \times 16.0)}$ = 18.073 / 18.07 / 18.1 / 18 (mol dm⁻³) TE on mass of H₂SO₄ in 1 dm³</p> <p>Allow 98 for molar mass of H₂SO₄ giving 18.092 / 18.09 / 18.1 / 18 (mol dm⁻³)</p> <p>Correct answer to 3 or more SF with no working scores (2)</p> <p>Do not award (2) for 18 unless 0.985 has been used in calculation</p>	(2)

Question Number	Answer	Additional Guidance	Mark
18(b)(ii)	<ul style="list-style-type: none"> calculation of $[H^+(aq)] / [H_3O^+(aq)]$ 	<p>Example of calculation:</p> $[H^+(aq)] / [H_3O^+(aq)] = 10^{-0.97}$ $= 0.10715 / 0.1072 / 0.107 / 0.11 \text{ (mol dm}^{-3}\text{)}$ <p>Ignore SF except 1 SF Ignore incorrect units</p> <p>Correct answer with no working scores (1) Do not award 0.1 / 0.10 / 0.214</p>	(1)

Question Number	Answer	Additional Guidance	Mark
18(b)(iii)	<p>An explanation that makes reference to the following points:</p> <p>First equilibrium</p> <ul style="list-style-type: none"> the first ionisation of sulfuric acid is complete or the equilibrium position of the first equation lies very far to the right <p>Second equilibrium</p> <ul style="list-style-type: none"> so $[H_3O^+(aq)]$ (from the second equilibrium) is very small 	<p>Allow $[H^+(aq)]$ for $[H_3O^+(aq)]$ Ignore missing state symbols</p> <p>Allow high $[H_3O^+(aq)]$ from first equilibrium Allow acid fully dissociates in first equilibrium (1) Ignore just acid fully dissociates</p> <p>(1) Allow second equilibrium shifts to the left Allow second dissociation is suppressed / further dissociation is prevented</p>	(2)

Question Number	Answer	Additional Guidance	Mark
18(c)(i)	<ul style="list-style-type: none"> <li data-bbox="427 639 1245 675">• $\text{HSO}_4^- + \text{OH}^- \rightarrow \text{SO}_4^{2-} + \text{H}_2\text{O}$ (1) <li data-bbox="427 756 1245 868">• $\text{SO}_4^{2-} + \text{H}^+ \rightarrow \text{HSO}_4^-$ or $\text{SO}_4^{2-} + \text{H}_3\text{O}^+ \rightarrow \text{HSO}_4^- + \text{H}_2\text{O}$ (1) 	<p data-bbox="1296 272 1704 304">Allow equations in either order</p> <p data-bbox="1296 347 1832 416">Allow \rightleftharpoons provided equations written in directions shown</p> <p data-bbox="1296 459 1787 491">Ignore state symbols even if incorrect</p> <p data-bbox="1296 534 1935 603">Penalise non-ionic equations once only e.g. using HCl and NaOH</p> <p data-bbox="1296 646 1697 758">Allow $\text{HSO}_4^- \rightarrow \text{SO}_4^{2-} + \text{H}^+$ and $\text{H}^+ + \text{OH}^- \rightarrow \text{H}_2\text{O}$ for M1</p>	(2)


Question Number	Answer	Additional Guidance	Mark
18(c)(ii)	<ul style="list-style-type: none"> <li data-bbox="439 309 1070 341">• calculation of the concentration of SO_4^{2-} ions <li data-bbox="439 496 1081 528">• calculation of the concentration of HSO_4^- ions <li data-bbox="439 719 707 751">• expression for K_a <li data-bbox="439 874 864 978">• re-arrangement of expression and calculation of $[\text{H}^+]$ <li data-bbox="439 1098 707 1129">• calculation of pH 	<p data-bbox="1270 272 1581 304">Example of calculation:</p> <p data-bbox="1270 304 1659 376">$[\text{SO}_4^{2-}] = \frac{25.0 \times 0.150}{1000} \times \frac{1000}{100}$ $= 0.0375 \text{ (mol dm}^{-3}\text{)}$</p> <p data-bbox="1270 416 1877 448">Allow mol $\text{SO}_4^{2-} = 0.00375 / 3.75 \times 10^{-3} \text{ (mol)}$</p> <p data-bbox="1270 488 1671 560">$[\text{HSO}_4^-] = \frac{75.0 \times 0.100}{1000} \times \frac{1000}{100}$ $= 0.075 \text{ (mol dm}^{-3}\text{)}$</p> <p data-bbox="1270 600 1861 632">Allow mol $\text{HSO}_4^- = 0.0075 / 7.5 \times 10^{-3} \text{ (mol)}$</p> <p data-bbox="1270 639 1883 671">Do not award this mark if subtraction then done</p> <p data-bbox="1270 711 1805 783">$K_a = \frac{[\text{H}^+][\text{SO}_4^{2-}]}{[\text{HSO}_4^-]} / 0.012 = \frac{[\text{H}^+] \times 0.0375}{0.075}$</p> <p data-bbox="1270 791 1854 823">Allow mol substituted into correct expression</p> <p data-bbox="1270 863 1720 935">$[\text{H}^+] = \frac{K_a[\text{HSO}_4^-]}{[\text{SO}_4^{2-}]} = \frac{0.012 \times 0.075}{0.0375}$</p> <p data-bbox="1270 967 1585 999">$= 0.024 \text{ (mol dm}^{-3}\text{)}$</p> <p data-bbox="1270 1007 1872 1038">TE on expression, $[\text{SO}_4^{2-}]$ and $[\text{HSO}_4^-]$ or mol</p> <p data-bbox="1270 1078 1659 1150">$\text{pH} = -\log[\text{H}^+] = -\log 0.024$ $= 1.6198 / 1.620 / 1.62 / 1.6$</p> <p data-bbox="1270 1158 1413 1190">TE on $[\text{H}^+]$</p> <p data-bbox="1270 1230 1816 1302">Ignore SF except 1 SF Correct answer without working scores (5)</p> <p data-bbox="1270 1342 1615 1374">Allow alternative methods</p>	(5)

(Total for Question 18 = 20 marks)













Question Number	Answer	Additional Guidance	Mark
19(a)	<ul style="list-style-type: none"><li data-bbox="383 309 539 336">• C₁₀H₁₈O	Allow symbols in any order i.e. C ₁₀ OH ₁₈ / H ₁₈ C ₁₀ O / H ₁₈ OC ₁₀ / OC ₁₀ H ₁₈ / OH ₁₈ C ₁₀ Allow large numbers e.g. C10H18O Do not award superscripts e.g. C ¹⁰ H ¹⁸ O	(1)


Question Number	Answer	Additional Guidance	Mark
19(b)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> • curly arrow from lone pair on C of CN⁻ towards C of aldehyde group (1) • curly arrow from C=O to, or just beyond, O and dipole on C=O (1) • intermediate (1) • curly arrow from lone pair on O⁻ to H and curly arrow from H-CN bond to anywhere on CN and final organic product (1) 	<p>Penalise an incorrect dipole in M1 and M4 once only Penalise curly arrow not starting from lone pair once only in M1 and M4 Penalise half arrow-heads once only</p> <p>Allow CN⁻ to attack C=O from any angle Allow CN bond displayed</p> <p>If M1 lost as curly arrow from N of CN⁻, allow CN joined to carbon through N Ignore connectivity for vertical CN groups</p> <p>Allow curly arrow from lone pair on O⁻ to H⁺ Ignore missing dipole in HCN</p>	(4)

Example of mechanism:



Question Number	Answer	Additional Guidance	Mark
19 (c)(i)	<ul style="list-style-type: none"> CH₃CO– / –COCH₃ / methyl ketone 	<p>Allow any combination of structural / displayed formula or skeletal formula</p> <p>Allow methyl next to ketone / methyl and ketone / methylcarbonyl</p> <p>Allow CH₃COR / RCOCH₃</p> <p>Ignore missing continuation bond from structures</p> <p>Do not award ethanal / methyl secondary alcohol / a specific compound</p>	(1)

Question Number	Answer	Additional Guidance	Mark				
19(c)(ii)	<ul style="list-style-type: none"> any 2 skeletal formulae (1) remaining 2 skeletal formulae (1) 	<p>Examples of skeletal formulae:</p> <table border="1" style="width: 100%; text-align: center;"> <tbody> <tr> <td></td> <td></td> </tr> <tr> <td></td> <td></td> </tr> </tbody> </table> <p>Ignore bond lengths and bond angles</p> <p>Allow (1) for 4 correct displayed / structural formulae</p>					(2)
							
							

Question Number	Answer	Additional Guidance	Mark
19(c)(iii)	<ul style="list-style-type: none"> • displayed formula of F • carbon atoms labelled 	<p>Example of displayed formula:</p>  <p>(1) Allow CH₃</p> <p>(1) Allow other unambiguous labels for the carbon atoms</p> <p>Allow M2 for labels on structural / skeletal formulae, including labels on formulae in (c)(ii)</p> <p>Ignore reference to singlet / splitting patterns</p>	(2)


Question Number	Answer	Additional Guidance	Mark												
19(d)*	<p>This question assesses a student's ability to show a coherent and logically structured answer with linkages and fully-sustained reasoning.</p> <p>Marks are awarded for indicative content and for how the answer is structured and shows lines of reasoning.</p> <p>The following table shows how the marks should be awarded for indicative content.</p> <table border="1" data-bbox="320 635 835 978"> <thead> <tr> <th>Number of indicative marking points seen in answer</th> <th>Number of marks awarded for indicative marking points</th> </tr> </thead> <tbody> <tr> <td>6</td> <td>4</td> </tr> <tr> <td>5-4</td> <td>3</td> </tr> <tr> <td>3-2</td> <td>2</td> </tr> <tr> <td>1</td> <td>1</td> </tr> <tr> <td>0</td> <td>0</td> </tr> </tbody> </table> <p>The following table shows how the marks should be awarded for structure and lines of reasoning.</p>	Number of indicative marking points seen in answer	Number of marks awarded for indicative marking points	6	4	5-4	3	3-2	2	1	1	0	0	<p>Guidance on how the mark scheme should be applied: The mark for indicative content should be added to the mark for lines of reasoning. For example, an answer with five indicative marking points that is partially structured with some linkages and lines of reasoning scores 4 marks (3 marks for indicative content and 1 mark for partial structure and some linkages and lines of reasoning). If there are no linkages between points, the same five indicative marking points would yield an overall score of 3 marks (3 marks for indicative content and no marks for linkages).</p>	(6)
Number of indicative marking points seen in answer	Number of marks awarded for indicative marking points														
6	4														
5-4	3														
3-2	2														
1	1														
0	0														

		Number of marks awarded for structure of answer and sustained line of reasoning	<p>In general it would be expected that 5 or 6 indicative points would get 2 reasoning marks, and 3 or 4 indicative points would get 1 mark for reasoning, and 0, 1 or 2 indicative points would score zero marks for reasoning.</p> <p>Additional incorrect chemistry loses a structure and lines of reasoning mark (if 1 or 2 have been awarded).</p>	
Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning demonstrated throughout.	2			
Answer is partially structured with some linkages and lines of reasoning.	1			
Answer has no linkages between points and is unstructured.	0			
<p>Comment: Look for the indicative marking points first, then consider the mark for structure of answer and sustained line of reasoning</p>				

	<p>Indicative content</p> <ul style="list-style-type: none"> • IP1 London forces Pentane (only) has London forces / all have London forces • IP2 Butanal Butanal (also) has (permanent) dipole-dipole interactions • IP3 Propanoic acid Propanoic acid (also) has (dipole-dipole and) hydrogen bonding • IP4 Intermolecular forces The London forces have about the same strength as they have a similar number of electrons / pentane has 42, butanal has 40 (and propanoic acid has 40 electrons) or butanal has dipole-dipole interactions because it is polar / has a dipole on C=O or propanoic acid has hydrogen bonding because it contains OH / COOH group • IP5 Butanal and pentane Dipole-dipole interactions are stronger than London forces or more energy is needed to overcome the dipole-dipole interactions than London forces • IP6 Butanal and propanoic acid Hydrogen bonding is stronger than dipole-dipole interactions or more energy is needed to overcome hydrogen bonding than dipole-dipole interactions 	<p>Allow dispersion forces / van der Waals' forces / forces between an instantaneous dipole and an induced dipole for London forces throughout answer Do not award IP1 if any other forces mentioned</p> <p>Allow dipole-dipole forces / attractions / bonds Do not award IP2 if hydrogen bonding included</p> <p>Allow similar relative molecular masses (butanal 72, pentane 72, propanoic acid 74) Do not award incorrect numbers of electrons / relative molecular masses</p> <p>Allow dipole-dipole interactions linked to C=O</p> <p>Allow diagram of hydrogen bond between two molecules Ignore formation of dimer</p> <p>Allow London forces are the weakest (intermolecular force) Do not award if covalent bonds broken or explanation is about intermolecular forces with water</p> <p>Allow hydrogen bonding is the strongest (intermolecular force) Do not award if covalent bonds broken or explanation is about intermolecular forces with water</p>	<p>(6)</p>
--	---	---	-------------------


(Total for Question 19 = 16 marks)

Question Number	Answer	Additional Guidance	Mark
20(a)	<ul style="list-style-type: none"> • substitution of values into expression for ΔS_{system} (1) • calculation of ΔS_{system} (1) • substitution of values into expression for $\Delta S_{\text{surroundings}}$ (1) • calculation of $\Delta S_{\text{surroundings}}$ (1) • calculation of ΔS_{total} (1) 	<p>Example of calculation:</p> $\Delta S_{\text{system}} = (2 \times 95.9) + (3 \times 205.0) - (2 \times 149.2)$ <p>Allow $191.8 + 615.0 - 298.4 / 806.8 - 298.4$</p> $\Delta S_{\text{system}} = (+)508.4 \text{ (J K}^{-1} \text{ mol}^{-1}\text{)}$ $\Delta S_{\text{surroundings}} = - \frac{(-67.2)}{298}$ $\Delta S_{\text{surroundings}} = (+)0.2255 \text{ (kJ K}^{-1} \text{ mol}^{-1}\text{)}$ <p>or $(+)225.5 \text{ (J K}^{-1} \text{ mol}^{-1}\text{)}$</p> $\Delta S_{\text{total}} = 508.4 + 225.5 = (+)733.9 \text{ (J K}^{-1} \text{ mol}^{-1}\text{)}$ <p>or $(+)0.7339 \text{ (kJ K}^{-1} \text{ mol}^{-1}\text{)}$</p> <p>TE on calculated values for ΔS_{system} and $\Delta S_{\text{surroundings}}$ Do not award ΔS_{system} added to $\Delta S_{\text{surroundings}}$ in different units e.g. $508.4 + 0.2255 = 508.6255$</p> <p>Ignore SF except 1 SF</p> <p>Units are not needed. Units in any order e.g. $\text{J mol}^{-1} \text{ K}^{-1}$ Penalise incorrect / incomplete units in ΔS_{total} but allow: e.g. J / mol/ K or J/mol.K mol^{-1} and K^{-1} in otherwise correct units</p> <p>Correct answer with no working scores (5)</p>	(5)

Question Number	Answer	Additional Guidance	Mark
20(b)(i)	<ul style="list-style-type: none"> • working for at least one half-life shown on graph • values of two half-lives • first order and because the half-lives are constant / the same / similar 	<p>Example of half-lives:</p>  <p>(1)</p> <p>(1)</p> <p>half-lives may be written or clearly shown on graph first half-life = 200 s and second half-life = 200 s ± 20 s Allow 2nd half-life = 400 – 200 = 200 s</p> <p>Stand alone mark</p> <p>Do not award zero order / second order</p> <p>(1)</p>	(3)

Question Number	Answer	Additional Guidance	Mark
20(b)(ii)	<p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> • first order with respect to BrO_3^- ions and because in runs 1 and 2 as $[\text{BrO}_3^-]$ doubles (and $[\text{H}^+]$ is constant) the rate doubles • in runs 1 and 3, as $[\text{BrO}_3^-]$ triples rate should triple (to 1.08×10^{-2}) and $[\text{H}^+]$ also doubles and rate increases by a factor of 4 • so reaction is second order with respect to H^+ ions 	<p>In M1 and M2, the working may be shown in the table Allow implied runs e.g. as $[\text{BrO}_3^-]$ doubles (and $[\text{H}^+]$ is constant), the rate doubles</p> <p>(1)</p> <p>Allow correct alternative explanations using runs 2 and 3 and others e.g. $[\text{BrO}_3^-]$ triples, $[\text{H}^+]$ doubles and rate x 12 Do not award just $[\text{H}^+]$ doubles and rate x 4 with no mention of bromate ions</p> <p>(1)</p> <p>Stand alone mark</p>	(3)

Question Number	Answer	Additional Guidance	Mark
20(b)(iii)	<ul style="list-style-type: none"> • rate equation • units of k 	<p>(1) rate = $k[\text{Br}^-(\text{aq})][\text{BrO}_3^-(\text{aq})][\text{H}^+(\text{aq})]^2$ TE on orders in (a)(i) and (ii) Allow species in any order / R for rate / K for k Ignore missing state symbols If no order given in (a)(i), allow rate equation with Br^- included or omitted</p> <p>(1) $\text{dm}^9 \text{mol}^{-3} \text{s}^{-1}$ Allow these in any order Allow $\text{dm}^9 \text{mol}^{-3} \text{s}^{-}$</p> <p>TE on rate equation</p>	(2)

Question Number	Answer	Additional Guidance	Mark
20(c)		<p data-bbox="1012 271 1254 303">Example of graph:</p>  <p data-bbox="1012 989 1288 1029">Example of gradient:</p> $\frac{-9.8 - (-5.85)}{3.3 \times 10^{-3} - 3.0 \times 10^{-3}} = -13167 \text{ K}$	(7)

Question Number	Answer	Additional Guidance	Mark
20(c)	<ul style="list-style-type: none"> <li data-bbox="416 272 981 379">• axes correct way around and suitable scale (1) <li data-bbox="416 459 981 566">• both axes labelled and units for x axis (1) <li data-bbox="416 646 981 753">• all points plotted correctly and best-fit straight line (1) <li data-bbox="416 801 981 826">• calculation of gradient (1) <li data-bbox="416 954 981 979">• sign and units of gradient (1) <li data-bbox="416 1107 981 1165">• calculation of activation energy (1) <li data-bbox="416 1257 981 1342">• sign and corresponding units of activation energy (1) 	<p data-bbox="1021 272 1821 416">Points / line must cover at least half the grid in both directions ln <i>k</i> values must become more negative down the axis with negative signs shown Allow horizontal axis shown at bottom of graph</p> <p data-bbox="1021 459 1865 596">y axis: ln <i>k</i> with no units on y axis and x axis: 0.0033 etc with (1/T) / K⁻¹ or 3.3 etc with (1/T) / 10⁻³ K⁻¹ or 3.3 x 10⁻³ etc with (1/T) / K⁻¹ or 3.3 etc with (1/T) x 10³ / K⁻¹ Brackets are not needed around 1/T</p> <p data-bbox="1021 646 1648 746">Allow ± ½ square Allow line covering points provided it is straight (1) Ignore extrapolation in either direction</p> <p data-bbox="1021 794 1659 932">This may be shown on the graph Allow gradient in the range (-)12800 to (-)13800 Allow gradient calculated from data in the table If gradient not evaluated, allow correct working</p> <p data-bbox="1021 948 1547 1048">(1) Negative sign and units K Allow 1/K⁻¹ for units Allow -12.8 to -13.8 kK for M4 and M5</p> <p data-bbox="1021 1091 1487 1228">$E_a = 13167 \times 8.31 / 1000 = 109.418$ Expected range 106 to 115 or $13167 \times 8.31 = 109418$ TE on gradient</p> <p data-bbox="1021 1273 1653 1410">+ 109.418 kJ mol⁻¹ or +109418 J mol⁻¹ (1) Allow kJ mol⁻¹ or J mol⁻¹ Ignore missing + but do not award - sign Penalise 1 SF for gradient and <i>E_a</i> value once only</p>	

(Total for Question 20 = 20 marks)

