

**CAMBRIDGE INTERNATIONAL EXAMINATIONS**

Cambridge International General Certificate of Secondary Education

## **MARK SCHEME for the March 2015 series**

### **0620 CHEMISTRY**

**0620/32**

Paper 3 (Extended Theory), maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the March 2015 series for most Cambridge IGCSE® components.

® IGCSE is the registered trademark of Cambridge International Examinations.

<b>Page 2</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>Cambridge IGCSE – March 2015</b>	<b>0620</b>	<b>32</b>

- 1 (a) chlorine/argon [1]
- (b) chlorine [1]
- (c) magnesium [1]
- (d) argon [1]
- (e) aluminium [1]
- (f) sodium [1]

[Total:6]

- 2 (a) Atoms of the same element/ atoms with same proton number/ atoms with same atomic number [1]
- different neutron number/ nucleon number/ mass number [1]

(b)

particle	number of protons	number of electrons	number of neutrons	nucleon number	symbol or formula
A					
B				23 (1)	Na(1) <sup>+</sup> (1)
C		10(1)		16(1)	
D	13 (1)		15 (1)		

[7]

[Total:9]

Page 3	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – March 2015	0620	32

- 3 (a) (making) fertilisers / nitric acid / nylon / explosives / urea  
(for) cleaning products (allow oven cleaner) / refrigeration [1]
- (b) equilibrium / reversible [1]
- (c) (nitrogen) air / atmosphere [1]  
  
(hydrogen) methane / water / steam / alkane / named alkane / hydrocarbon / crude oil  
or petroleum / natural gas [1]
- (d) iron [1]
- (e) (i) rate increases / faster [1]  
More (effective) collisions [1]  
(ii) yield decreases [1]  
(forward reaction) exothermic / reverse reaction endothermic / high temp  
favours endothermic reaction [1]
- (f) (i) yield increases [1]  
  
less / fewer molecules or moles or volume on RHS OR / high pressure  
favours reaction which produces fewer molecules or moles or volume [1]  
(ii) particles / molecules closer / more particles per unit area or volume / more  
molecules per unit area or volume / more concentration / particles have less  
space between them **and** more collisions [1]  
(iii) safety issues / higher cost [1]
- (g) 3 bond pairs between N & H [1]  
  
Lone pair on N [1]
- (h) (i) proton / H<sup>+</sup> acceptor [1]  
(ii)  $2\text{NH}_3 + \text{H}_2\text{SO}_4 \rightarrow (\text{NH}_4)_2\text{SO}_4$  [2]  
  
Formula of (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> (1)  
The rest (1)

[Total:18]

Page 4	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – March 2015	0620	32

- 4 (a) (i) 82.76/12 and 17.2(4)/(1)  
or evaluation: 6.89 / 6.9(0) and 17.2(4) [1]
- C<sub>2</sub>H<sub>5</sub> [1]
- OR**  
82.76/100 × 58 = 48 and 17.24/100 × 58 = 10  
or evaluation i.e. 48 and 10 [1]
- C<sub>2</sub>H<sub>5</sub> [1]
- (ii) (C<sub>2</sub>H<sub>5</sub> =) 29 [1]
- (58/29 = 2) C<sub>4</sub>H<sub>10</sub> [1]
- OR:  
82.76/100 × 58 = 48 and 17.24/100 × 58 = 10  
or evaluation i.e. 48 and 10 [1]
- 48/12 = 4 10/1 = 10 (therefore) C<sub>4</sub>H<sub>10</sub> [1]
- (b) (i) C<sub>n</sub>H<sub>2n</sub> [1]
- (ii) CH<sub>2</sub> [1]
- (c) (contains) double bond / triple bond / multiple bond(s) / not all bonds are single [1]
- (contains) carbon and hydrogen **only** [1]
- (d) bromine / bromine water [1]
- no change / stays brown / orange / yellow / red-brown or only changes in UV [1]
- (brown / orange / yellow) to colourless / decolourised [1]
- (e) (i) circle / brackets around any 2 consecutive carbon atoms in the main chain and all attached atoms [1]
- e.g.
- The diagram shows the structural formula of butane, CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>. The first two carbon atoms and their attached groups (C<sub>2</sub>H<sub>5</sub> and H) are enclosed in a rectangular box. The structure is drawn as a horizontal chain of four carbon atoms, each with its bonds to other atoms shown as vertical lines. The first carbon is bonded to a C<sub>2</sub>H<sub>5</sub> group and a hydrogen atom. The second carbon is bonded to a hydrogen atom. The third carbon is bonded to a C<sub>2</sub>H<sub>5</sub> group and a hydrogen atom. The fourth carbon is bonded to a hydrogen atom. The chain continues to the left and right of the first and last carbons, indicated by horizontal lines.
- (ii) CH<sub>3</sub>CH<sub>2</sub>CH=CH<sub>2</sub> / C<sub>2</sub>H<sub>5</sub>CH=CH<sub>2</sub> (double bond must be shown) [1]
- butene / but-1-ene [1]

Page 5	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – March 2015	0620	32

(iii)  $(\text{CH}_3)_2\text{C}=\text{CH}_2 / \text{CH}_3\text{CH}=\text{CHCH}_3 / (\text{CH}_2)_2\text{CHCH}_3 / (\text{CH}_2)_4$  [1]

[Total:15]

5 (a) Bauxite [1]

(b) carbon/graphite [1]

(c) improves conductivity/better conductor [1]

Lower (operating) temperature/save energy/saves electricity/saves heat [1]

(d) anode:  $2\text{O}^{2-} \rightarrow \text{O}_2 + 4\text{e}^- / 2\text{O}^{2-} - 4\text{e}^- \rightarrow \text{O}_2$  [1]

cathode:  $\text{Al}^{3+} + 3\text{e}^- \rightarrow \text{Al} / \text{Al}^{3+} \rightarrow \text{Al} - 3\text{e}^-$  [1]

(e) (i) Iron carbon aluminium/Fe, C, Al [1]

(ii) Aluminium oxide is not reduced by carbon but iron(III) oxide is [1]

(f) haematite/hematite [1]

(g) **Allow:** multiples in (i) to (iv)

(i)  $\text{C} + \text{O}_2 \rightarrow \text{CO}_2$  [1]

(ii)  $\text{CO}_2 + \text{C} \rightarrow 2\text{CO}$  [1]

(iii)  $\text{Fe}_2\text{O}_3 + 3\text{CO} \rightarrow 2\text{Fe} + 3\text{CO}_2 / \text{Fe}_2\text{O}_3 + 3\text{C} \rightarrow 2\text{Fe} + 3\text{CO} /$   
 $2\text{Fe}_2\text{O}_3 + 3\text{C} \rightarrow 4\text{Fe} + 3\text{CO}_2$  [1]

(iv)  $\text{CaO} + \text{SiO}_2 \rightarrow \text{CaSiO}_3 / \text{CaCO}_3 + \text{SiO}_2 \rightarrow \text{CaSiO}_3 + \text{CO}_2$  [1]

[Total:13]

6 (a) Any **two** from:

- bubbles/effervescence/fizzing
- (some of the) solid/copper carbonate dissolves/disappears **or** some (brown) solid seen (undissolved)
- (colourless) solution or liquid turns blue [2]

Page 6	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – March 2015	0620	32

- (b) filter / centrifuge / decant [1]  
wash with (distilled) water [1]  
(dry with) filter paper / tissues / warm windowsill / in sun / oven / fan / heat [1]

- (c) (i) Blue precipitate / ppt [1]  
(ii)  $\text{Cu}^{2+} + 2\text{OH}^- \rightarrow \text{Cu}(\text{OH})_2$  [1]

- (d) (i)  $\text{Cu}(\text{OH})_2(\text{s}) \rightarrow \text{CuO}(\text{s}) + \text{H}_2\text{O}(\text{g})$   
Equation [1]  
State symbols of correct chemical equation [1]  
(ii) carbon / hydrogen [1]

[Total:10]

- 7 (a) Any **two** from:  
yeast / 20–40 °C / anaerobic or without oxygen or without air / (aqueous)  
solution or water or aqueous [2]

- (b) (i)  $M_r = 180 (1) (30/180) = 0.167 (1)$  [2]  
(ii)  $2 \times 0.167$  or  $2 \times 46$  or  $0.333$  or  $92$  [1]  
 $(2 \times 0.167 \times 46) = 15.3(33) (\text{g})$  [1]  
(iii)  $(2 \times 0.167 \times 24) = 8 (\text{dm}^3)$  [1]

- (c) (i) Crude oil / petroleum [1]  
(ii)  $\text{C}_2\text{H}_4 + \text{H}_2\text{O} \rightarrow \text{C}_2\text{H}_5\text{OH} / \text{CH}_3\text{CH}_2\text{OH}$  [1]

[Total:9]