



UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS  
International General Certificate of Secondary Education

CANDIDATE  
NAME

CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--



**CHEMISTRY**

**0620/31**

Paper 3 (Extended)

**May/June 2013**

**1 hour 15 minutes**

Candidates answer on the Question Paper.

No Additional Materials are required.

**READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

**DO NOT WRITE IN ANY BARCODES.**

Answer **all** questions.

Electronic calculators may be used.

A copy of the Periodic Table is printed on page 12.

You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

This document consists of **12** printed pages.



1 Petroleum contains hydrocarbons which are separated by fractional distillation.

(a) (i) Complete the following definition of a hydrocarbon.

A hydrocarbon is a compound which .....  
..... [2]

(ii) Explain what is meant by the term *fractional distillation*.

.....  
.....  
..... [2]

(b) Some of the fractions obtained from petroleum are given below.  
State a use for each fraction.

bitumen .....

lubricating fraction .....

paraffin fraction .....

gasoline fraction ..... [4]

[Total: 8]

2 An element, **M**, has the electron distribution 2 + 8 + 18 + 3.

(a) Which group in the Periodic Table is element **M** likely to be in?

..... [1]

(b) Predict whether element **M** is a poor or a good conductor of electricity.  
Give a reason for your answer.

..... [1]

(c) Binary compounds contain two atoms per molecule, for example  $\text{HCl}$ .  
Identify an element which could form a binary compound with element **M**.

..... [1]

(d) Predict the formula of the sulfate of **M**. The formula of the sulfate ion is  $\text{SO}_4^{2-}$ .

..... [1]

- (e) The hydroxide of **M** is a white powder which is insoluble in water. Describe how you could show that this hydroxide is amphoteric.

.....  
 .....  
 ..... [2]

[Total: 6]

- 3 A small piece of marble,  $\text{CaCO}_3$ , was added to  $5.0 \text{ cm}^3$  of hydrochloric acid, concentration  $1.0 \text{ mol/dm}^3$ , at  $25^\circ\text{C}$ . The time taken for the reaction to stop was measured. The experiment was repeated using  $5.0 \text{ cm}^3$  of different solutions of acids. The acid was in excess in all of the experiments.

Typical results are given in the table.

experiment	temperature/ $^\circ\text{C}$	acid solution	time/min
1	25	hydrochloric acid $1.0 \text{ mol/dm}^3$	3
2	25	hydrochloric acid $0.5 \text{ mol/dm}^3$	7
3	25	ethanoic acid $1.0 \text{ mol/dm}^3$	10
4	15	hydrochloric acid $1.0 \text{ mol/dm}^3$	8

- (a) (i) Explain why it is important that the pieces of marble are the same size and the same shape.

.....  
 .....  
 ..... [2]

- (ii) How would you know when the reaction had stopped?

..... [1]

- (b) The equation for the reaction in experiment 1 is:



Complete the following ionic equation.



[1]

- (c) (i) Explain why the reaction in experiment 1 is faster than the reaction in experiment 2.

.....  
 ..... [1]

- (ii) The acids used for experiment 1 and experiment 3 have the same concentration. Explain why experiment 3 is slower than experiment 1.


.....  
 .....  
 ..... [2]

- (iii) Explain in terms of collisions between reacting particles why experiment 4 is slower than experiment 1.

.....  
 .....  
 ..... [3]

[Total: 10]

- 4 The structural formula of cyclohexane is drawn below.



- (a) The name gives information about the structure of the compound. **Hex** because there are six carbon atoms and **cyclo** because they are joined in a ring. What information about the structure of this compound is given by the ending **ane**?

.....  
 ..... [2]

- (b) What are the molecular and empirical formulae of cyclohexane?

molecular formula .....

empirical formula .....

[2]

(c) Draw the structural formula of cyclobutane.

[1]

(d) (i) Deduce the molecular formula of hexene.

..... [1]

(ii) Explain why cyclohexane and the alkene, hexene, are isomers.

.....

.....

..... [2]

(e) Describe a test which would distinguish between cyclohexane and the unsaturated hydrocarbon hexene.

test .....

result of test with cyclohexane .....

.....


result of test with hexene .....

..... [3]

[Total: 11]

5 The reactivity series shows the metals in order of reactivity.

- (a) The reactivity series can be established using displacement reactions. A piece of zinc is added to aqueous lead nitrate. The zinc becomes coated with a black deposit of lead.



Zinc is more reactive than lead.

The reactivity series can be written as a list of ionic equations.

.....  $\rightarrow$  ..... + ..... most reactive metal: the best reductant (reducing agent)



- (i) In the space at the top of the list, write an ionic equation for a metal which is more reactive than zinc. [1]

- (ii) Write an ionic equation for the reaction between aqueous silver(I) nitrate and zinc.

..... [2]

- (iii) Explain why the positive ions are likely to be oxidants (oxidising agents).

..... [1]


- (iv) Deduce which ion is the best oxidant (oxidising agent).

..... [1]

- (v) Which ion(s) in the list can oxidise lead metal?

..... [1]

- (b) A reactivity series can also be established by measuring the voltage of simple cells. The diagram shows a simple cell.



Results from cells using the metals tin, cadmium, zinc and copper are given in the table below.

cell	electrode 1 positive electrode	electrode 2 negative electrode	voltage /volts
1	copper	cadmium	0.74
2	copper	tin	0.48
3	copper	zinc	1.10

Write the four metals in order of increasing reactivity and explain how you used the data in the table to determine this order.

.....  
 .....  
 ..... [3]

[Total: 9]

- 6 Ammonia is a compound which only contains the elements nitrogen and hydrogen. It is a weak base.

- (a) (i) Define the term *base*.

..... [1]


- (ii) Given aqueous solutions of ammonia and sodium hydroxide, both having a concentration of  $0.1 \text{ mol/dm}^3$ , how could you show that ammonia is the weaker base?

.....  
 .....  
 ..... [2]

- (b)** Ammonia is manufactured by the Haber Process. The economics of this process require that as much ammonia as possible is made as quickly as possible. Explain how this can be done using the following information.

The conditions for the following reversible reaction are:

- 450 °C
- 200 atmospheres pressure
- iron catalyst



.....

.....

.....

.....

.....

.....

.....


..... [5]

- (c)** Another compound which contains only nitrogen and hydrogen is hydrazine,  $\text{N}_2\text{H}_4$ .

Complete the equation for the preparation of hydrazine from ammonia.



- (d)** The structural formula of hydrazine is given below.



Draw a diagram showing the arrangement of the valency electrons in one molecule of the covalent compound hydrazine.

Use x to represent an electron from a nitrogen atom.

Use o to represent an electron from a hydrogen atom.

[3]



(e) Hydrazine is a weak base and it removes dissolved oxygen from water. It is added to water in steel boilers to prevent rusting.

(i) One way it reduces the rate of rusting is by changing the pH of water. What effect would hydrazine have on the pH of water?

..... [1]


(ii) Give a reason, other than pH, why hydrazine reduces the rate of rusting.

..... [1]

[Total: 15]

7 The hydroxides of the Group I metals are soluble in water. Most other metal hydroxides are insoluble in water.

(a) (i) Crystals of lithium chloride can be prepared from lithium hydroxide by titration.



25.0 cm<sup>3</sup> of aqueous lithium hydroxide is pipetted into the conical flask. A few drops of an indicator are added. Dilute hydrochloric acid is added slowly to the alkali until the indicator just changes colour. The volume of acid needed to neutralise the lithium hydroxide is noted.


A neutral solution of lithium chloride, which still contains the indicator, is left. Describe how you could obtain a neutral solution of lithium chloride which does **not** contain an indicator.

.....  
..... [2]

- (ii) You cannot prepare a neutral solution of magnesium chloride by the same method. Describe how you could prepare a neutral solution of magnesium chloride.

.....  
 .....  
 ..... [3]

- (b) The concentration of the hydrochloric acid was  $2.20 \text{ mol/dm}^3$ . The volume of acid needed to neutralise the  $25.0 \text{ cm}^3$  of lithium hydroxide was  $20.0 \text{ cm}^3$ . Calculate the concentration of the aqueous lithium hydroxide.



.....  
 .....  
 ..... [2]

- (c) Lithium chloride forms three hydrates. They are  $\text{LiCl}\cdot\text{H}_2\text{O}$ ,  $\text{LiCl}\cdot 2\text{H}_2\text{O}$  and  $\text{LiCl}\cdot 3\text{H}_2\text{O}$ . Which **one** of these three hydrates contains 45.9% of water? Show how you arrived at your answer.

.....  
 .....  
 ..... [3]

[Total: 10]

- 8 There are three types of giant structure - ionic, metallic and giant covalent.

- (a) In an ionic compound, the ions are held in a lattice by strong forces.

- (i) Explain the term *lattice*.

.....  
 ..... [2]

- (ii) Explain how the ions are held together by strong forces.

.....  
 ..... [1]

(b) Describe the bonding in a typical metal.

.....  
 .....  
 ..... [3]

(c) The electrical conductivities of the three types of giant structure are given in the following table.

type of structure	conductivity of solid	conductivity of liquid
ionic	poor	good
metallic	good	good
giant covalent	poor	poor

Explain the differences in electrical conductivity between the three types of giant structure and the difference, if any, between the solid and liquid states of the same structure.

.....  
 .....  
 .....  
 .....  
 ..... [5]

[Total: 11]

**DATA SHEET**  
**The Periodic Table of the Elements**

Group		I	II	III	IV	V	VI	VII	0							
		1 <b>H</b> Hydrogen 1							2 <b>He</b> Helium 2							
3	4	7 <b>Li</b> Lithium	9 <b>Be</b> Beryllium		11 <b>B</b> Boron	12 <b>C</b> Carbon	13 <b>Al</b> Aluminium	14 <b>Si</b> Silicon	15 <b>P</b> Phosphorus	16 <b>S</b> Sulfur	17 <b>Cl</b> Chlorine	18 <b>Ar</b> Argon				
11	12	23 <b>Na</b> Sodium	24 <b>Mg</b> Magnesium		27 <b>Fe</b> Iron	28 <b>Ni</b> Nickel	29 <b>Cu</b> Copper	30 <b>Zn</b> Zinc	31 <b>Ga</b> Gallium	32 <b>Ge</b> Germanium	33 <b>As</b> Arsenic	34 <b>Se</b> Selenium	35 <b>Br</b> Bromine	36 <b>Kr</b> Krypton		
19	20	39 <b>K</b> Potassium	40 <b>Ca</b> Calcium		44 <b>Ru</b> Ruthenium	45 <b>Rh</b> Rhodium	46 <b>Pd</b> Palladium	47 <b>Ag</b> Silver	48 <b>Cd</b> Cadmium	49 <b>In</b> Indium	50 <b>Sn</b> Tin	51 <b>Sb</b> Antimony	52 <b>Te</b> Tellurium	53 <b>I</b> Iodine	54 <b>Xe</b> Xenon	
37	38	85 <b>Rb</b> Rubidium	88 <b>Sr</b> Strontium		101 <b>Ru</b> Ruthenium	102 <b>Rh</b> Rhodium	103 <b>Pd</b> Palladium	104 <b>Ag</b> Silver	105 <b>Cd</b> Cadmium	106 <b>In</b> Indium	107 <b>Sn</b> Tin	108 <b>Sb</b> Antimony	109 <b>Te</b> Tellurium	110 <b>I</b> Iodine	111 <b>Xe</b> Xenon	
55	56	133 <b>Cs</b> Caesium	137 <b>Ba</b> Barium		186 <b>Re</b> Rhenium	187 <b>Rh</b> Rhodium	188 <b>Pt</b> Platinum	189 <b>Au</b> Gold	190 <b>Hg</b> Mercury	191 <b>Tl</b> Thallium	192 <b>Pb</b> Lead	193 <b>Bi</b> Bismuth	194 <b>Po</b> Polonium	195 <b>At</b> Astatine	196 <b>Rn</b> Radon	
87	88	226 <b>Fr</b> Francium	226 <b>Ra</b> Radium		227 <b>Ac</b> Actinium	227 <b>Ac</b> Actinium	227 <b>Ac</b> Actinium	227 <b>Ac</b> Actinium	227 <b>Ac</b> Actinium	227 <b>Ac</b> Actinium	227 <b>Ac</b> Actinium	227 <b>Ac</b> Actinium	227 <b>Ac</b> Actinium	227 <b>Ac</b> Actinium	227 <b>Ac</b> Actinium	227 <b>Ac</b> Actinium

140 <b>Ce</b> Cerium	141 <b>Pr</b> Praseodymium	144 <b>Nd</b> Neodymium	150 <b>Sm</b> Samarium	152 <b>Eu</b> Europium	157 <b>Gd</b> Gadolinium	162 <b>Dy</b> Dysprosium	165 <b>Ho</b> Holmium	167 <b>Er</b> Erbium	169 <b>Tm</b> Thulium	173 <b>Yb</b> Ytterbium	175 <b>Lu</b> Lutetium		
58	59	60	62	63	64	66	67	68	69	70	71		
232 <b>Th</b> Thorium	232 <b>Pa</b> Protactinium	238 <b>U</b> Uranium	238 <b>Pu</b> Plutonium	238 <b>Np</b> Neptunium	238 <b>Am</b> Americium	238 <b>Cm</b> Curium	238 <b>Bk</b> Berkelium	238 <b>Cf</b> Californium	238 <b>Es</b> Einsteinium	238 <b>Fm</b> Fermium	238 <b>Md</b> Mendelevium	238 <b>No</b> Nobelium	238 <b>Lr</b> Lawrencium
90	91	92	94	95	96	98	97	98	99	100	101	102	103

a	X	b
Key	X	b

\*58-71 Lanthanoid series  
†90-103 Actinoid series

a = relative atomic mass  
X = atomic symbol  
b = proton (atomic) number

The volume of one mole of any gas is 24 dm<sup>3</sup> at room temperature and pressure (r.t.p.).

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.